

Draco 119: A Remarkable Heavy Element-deficient Giant¹

Jon. P. Fulbright

*Observatories of the Carnegie Institution of Washington, 813 Santa Barbara St., Pasadena,
CA 91101*

jfulb@ociw.edu

R. Michael Rich

*Division of Astronomy, Department of Physics and Astronomy, UCLA, Los Angeles, CA
90095-1562*

rmr@astro.ucla.edu

Sandra Castro

*Infrared Processing and Analysis Center, Caltech, 770 S. Wilson Blvd., Pasadena, CA
91125 and The European Southern Observatory, Karl-Schwarzschild-Str. 2, Garching,
D-85748*

smc@astro.caltech.edu

ABSTRACT

We report the abundance analysis of new high S/N spectra of the most metal-poor ($[\text{Fe}/\text{H}] = -2.95$) star presently known to be a member of a dwarf galaxy, the Draco dSph red giant, D119. No absorption lines for elements heavier than Ni are detected in two Keck HIRES spectra covering the $\lambda\lambda$ 3850–6655 Å wavelength range, phenomenon not previously noted in any other metal-poor star. We present upper limits for several heavy element abundances. The most stringent limits, based on the non-detection of Sr II and Ba II lines, indicate that the total s- and r-process enrichment of D119 is at least 100 times smaller than Galactic stars of similar metallicity. The light element abundances are consistent with the star having formed out of material enriched primarily by massive Type II supernovae ($M > 20\text{--}25 M_{\odot}$). If this is the case, we are forced to conclude that massive, metal-poor Type II supernovae did not contribute to the r-process in the proto-Draco environment. We compare the abundance pattern observed in D119 to current predictions of prompt enrichment and pair-instability supernovae and find that the model predictions fail by an order or magnitude or more for many elements.

Subject headings: stars: abundances; nuclear reactions, nucleosynthesis, abundances; galaxies: individual (Draco dSph)

1. Introduction

The elements heavier than the iron group are believed to be produced primarily by neutron-capture processes. In their landmark work on the origin of the elements, (Burbidge et al. 1957) introduced the idea that the production of heavy elements depends on whether the rate of neutron capture is fast relative to the beta-decay timescale of the resulting nucleus (“r-process”) or slow relative to beta-decay (“s-process”).

The exact astrophysical sites of these processes have been the basis of much research. Sneden & Cowan (2003) and Truran et al. (2002) review the current understanding of the origin of the neutron-capture elements. The s-process is generally accepted to occur in two sites: the He-burning cores of massive stars and the thermally pulsing He shells of AGB stars. However, the specific site of the r-process is an unsolved problem, but it is strongly suspected that Type II supernovae play an important role. Truran et al. (2002) point out four possible sites, three of which occur in Type II supernovae explosions.

The mass range of Type II supernova progenitors that host the r-process is also under debate. Mathews et al. (1992), Travaglio et al. (1999) and Wanajo et al. (2003) propose that 8–10 $M_{\odot}$ progenitors are the predominant contributors to the r-process. This mass range provides the best fit between their r-process production models and the observed distribution of neutron-capture elements as a function of [Fe/H]. Alternatively, Tsujimoto et al. (2000) suggest that the largest contributors to the r-process are 20–25 $M_{\odot}$ progenitors. They do not place an upper mass limit to r-process contributors, but they do not believe that the low mass Type II supernovae are significant contributors to the r-process.

Additionally, it has been suggested that an additional “prompt” enrichment episode is necessary to enrich the protogalactic medium before the formation of metal-poor stars. Qian & Wasserburg (2001), Qian & Wasserburg (2002) and Qian (2003) put forth a multi-component model for early chemical evolution. The three components of the model are a “prompt” (P) component that represents the contribution from very high mass ($M > 100 M_{\odot}$) Population III stars, plus “high”- and “low”-frequency (H and L) enrichment events,

¹Based on data obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA, and was made possible by the generous financial support of the W. M. Keck Foundation.

both identified with Type II supernova. The prediction for the prompt enrichment contains contributions of the light neutron-capture elements Sr, Y and Zr.

A possible source of any prompt enrichment episode may have been a population of zero-metallicity stars. Heger & Woosley (2002) and Umeda & Nomoto (2002) provide predictions of the yields of a first generation of very massive (100 to 300 $M_{\odot}$) stars. These stars explode as pair instability supernovae, and produce a distinctive nucleosynthesis pattern. These supernovae are not predicted to produce large quantities of the neutron-capture elements.

In this paper, we describe our analysis of the red giant star Draco 119 (D119; identification by Stetson 1979). The star is the most metal poor in the Keck/HIRES sample of Shetrone et al. (1998) which reports analysis of a relatively low S/N (~ 25) red spectrum of the giant. Their reported $[\text{Fe}/\text{H}]=-2.97$ placed the star in an interesting range of abundance: nearly 1 dex more metal poor than any Galactic globular cluster, and as metal poor as the point where the extreme Galactic halo shows striking new abundance trends (McWilliam et al. 1995; Ryan et al. 1996; Johnson 2002). The star was targeted for observation by one of us (RMR) to discover whether such a star would fit into the extrapolation of the abundance trends for Galactic globular cluster giants, or would rather show abundance patterns characteristic of the ultra metal-poor halo stars.

Even after detailed abundances have been analyzed for some 40 additional giants in every Milky Way dwarf spheroidal galaxy except for Leo II, D119 remains to date as the most metal-poor star presently documented in any Milky Way dwarf spheroidal galaxy (Shetrone et al. 1998; Bonifacio et al. 2000; Shetrone et al. 2001, 2003; McWilliam et al. 2003).

Compared to the effort of Shetrone et al. (1998) we have invested considerably more integration time and pushed blueward, yielding two spectra of considerably higher S/N. In the course of analyzing this star, we discovered the extreme deficiency of neutron-capture elements in this star, as immediately evidenced by the lack of a detectable Ba II 4554Å line, the first ever such case in a metal-poor cool giant.

Given the unique nature of D119, we compare it to the bright, well studied halo star HD 88609 which shares similar stellar parameters, and to BD +80 245, a low-alpha metal-poor halo star (Carney et al. 1997; Ivans et al. 2003) with low $[\text{Ba}/\text{Fe}]$, yet still not as extreme as D119. We finally compare the observed abundance distribution of D119 to predictions of early enrichment from a range of supernova models.

2. Observations and Abundance Analysis

The data are obtained using two settings of the HIRES spectrograph (Vogt et al. 1994) on the Keck I telescope. For both settings a slit returning a resolution of 45000 was used. The first setting, used on 6 July 2000, covers the wavelength range from $\lambda\lambda$ 3850–6280 Å. Five 3000 s exposures were taken with this setting. The second setting, used on 29 June 2001, covers the wavelength range of $\lambda\lambda$ 4225–6655 Å, with some gaps in the wavelength coverage. Seven 3000 s exposures were taken with this setting. The data were reduced using the MAKEE² data reduction package. The final signal-to-noise (S/N) varies as a function of wavelength, reaching over 60 per pixel in the red, but falling to ~ 5 per pixel in the blue. The S/N at the λ 4077 Å and λ 4215 Å Sr II lines is about 7 per pixel, while the S/N at the λ 4554 Å and λ 4934 Å Ba II lines is about 35 per pixel.

We checked the star for signs of binarity by comparing the observed heliocentric radial velocities. Matt Shetrone provided us with the 28 August 1997 HIRES spectrum used in Shetrone et al. (1998). The mean radial velocity for D119 was measured to be -293.9 ± 0.1 km/s. The difference in radial velocity between the July 2000 and June 2001 spectrum is -0.2 ± 0.1 km/s, while the difference in radial velocity between the August 1997 and June 2001 is $+0.2 \pm 0.3$ km/s. We do not detect any large radial velocity variations over the nearly four years spanned by the spectra.

The initial line list was taken from Johnson (2002), which studied similar metallicity giants in the Galactic halo. Additional lines were added from Johnson & Bolte (2002), Fulbright (2000) and sources therein. Hyperfine splitting was taken into account for Na I D, Al I, Mn I, Co I, Sr II, Ba II, and Eu II lines. The C abundance of the two stars were determined by CH lines in the G-band using a line list provided by Andy McWilliam based on Brown et al. (1987). The line list and equivalent width measurements for both stars are given in Table 1.

Many of the interesting absorption lines in D119 are very weak or undetectable in our spectra. To place meaningful upper limits on the abundances of these elements, it is necessary to calculate upper limits for the equivalent widths of these lines. The full Fe line list contains nearly 200 lines, but only 96 were used in the final abundance analysis. We reject many of the remaining lines because they are too weak to measure. The Fe abundance determined from the measured Fe lines can then be used to predict the equivalent widths of the weaker, unmeasured, Fe lines. By comparing the estimated equivalent widths of these lines against

²MAKEE was developed by T. A. Barlow specifically for reduction of Keck HIRES data. It is freely available on the world wide web at the Keck Observatory home page; <http://www2.keck.hawaii.edu:3636/>.

the observed spectrum, we estimate the detection limit as a function of wavelength and S/N.

The value of this limit is roughly twice that given by Cayrel (1988) for the measurement error of the equivalent width of a line. This “two-sigma limit” allows for the estimation of the measurement limit in regions where there are no Fe lines. The two-sigma limit is ~ 5 mÅ in the red, but increases to ~ 20 – 30 mÅ in the blue.

To compare D119 to similar stars in the Galactic halo, we have re-analyzed the well-studied halo star HD 88609. Previous analyses of HD 88609 (Johnson & Bolte 2002; Johnson 2002; Fulbright 2000) show that this star is typical of very metal-poor halo giants and has stellar parameters similar to those of D119. In the re-analysis, we adopt primarily the equivalent width measurements of HD 88609 from the three papers cited above. Additional lines were measured from the Keck/HIRES spectra kindly provided by Jennifer Johnson.

The abundance analysis is conducted using the LTE code MOOG (Snedden 1973) and Kurucz³ atmospheres that do not include overshooting. Using a model with overshooting raises most of the $\log \epsilon$ values of all the neutral species by ~ 0.1 dex, but has very little effect on the ionized species.

Shetrone et al. (1998) uses stellar parameters based on Fe lines, but Kraft & Ivans (2003) show that while T_{eff} values based on the excitation plot of Fe I may be reliable, surface gravities based on the ionization equilibrium of Fe are not reliable for metal-poor globular cluster giants. For this star, we adopt a T_{eff} value of 4440 K, derived from the Alonso et al. (1999a) B-V calibration, assuming a reddening to the Draco dSph of 0.03 mag (Mateo 1998). If a distance modulus of 19.58 to Draco is assumed (Mateo 1998), then D119 has $M_V = -2.2$. From Equation 1 of Kraft & Ivans (2003) and the bolometric corrections of Alonso et al. (1999a), the resulting $\log g$ value is 0.88. If the Shetrone et al. T_{eff} value of 4370 K is used, the $\log g$ value only drops to 0.86. This is considerably higher than the Shetrone et al. $\log g$ of 0.3. When our value of $\log g$ is used, the difference $\log n(\text{Fe I}) - \log n(\text{Fe II})$ is -0.02 dex.

We find a microturbulent velocity of 2.4 ± 0.1 km/s by forcing the abundance given by Fe I lines to be independent of line strength. The uncertainty in the microturbulent velocity, v_t , was computed by multiplying the uncertainty in the slope of the least-squares fit to the $\log \epsilon(\text{Fe I})$ vs. $\log(\text{Reduced Width})$ plot by the inverse of the change in this slope as a function of v_t .

For HD 88609, we adopt the Alonso et al. (1999b) T_{eff} value of 4600 K. Hanson et al. (1998) found $M_V = -1.2$ for this star, which results in a $\log g$ value of 1.38. We determine

³<http://kurucz.harvard.edu/>

a microturbulent velocity of 2.1 ± 0.1 km/s.

Uncertainties for the abundances are calculated assuming the errors in stellar parameters and equivalent widths are independent. The uncertainty from parameter errors are approximated by changing the T_{eff} value by 150 K and altering the $\log g$ and other parameters accordingly. The uncertainty from line measurements is based on the calculations discussed above. For D119, the dominant contribution to error in the abundance determination comes from errors in the equivalent widths, while for HD 88609, uncertainties in the stellar parameters are the dominant contribution to the error in the abundance determination.

The final abundances, uncertainties, and upper limits for D119 and HD 88609 are given in Table 2. As suggested by Kraft & Ivans (2003), we have used the Fe I abundance in determining the $[X/\text{Fe}]$ ratio from neutral species and the Fe II abundance for singly-ionized species and $[\text{O I}]$. One of the advantages of using this method for computing abundance ratios is that there is a reduced sensitivity to parameter changes.

3. D119 and the Early Enrichment

Strontium and barium have the strongest absorption lines for neutron-capture elements. In HD 88609, the $\lambda 4077$ Å and $\lambda 4215$ Å Sr II lines have equivalent widths of 167 mÅ and 160 mÅ and the $\lambda 4554$ Ba II line has an equivalent width of 91 mÅ. We did not detect these two lines, nor any lines of any element heavier than Ni in D119. Figure 1 shows spectral regions around the $\lambda 4215$ Å Sr II and $\lambda 4554$ Å Ba II lines for D119 and HD 88609.

The difference in Sr and Ba abundances in these two stars is clear from the figure, and is confirmed by the abundance analysis. The $[\text{Sr}/\text{Fe}]$ and $[\text{Ba}/\text{Fe}]$ abundance limits are about a factor of 100 less than what is seen in HD 88609, and the $[\text{Ba}/\text{Fe}]$ ratio is below the observed value for any metal-poor star. The upper limits of the $[\text{Y}/\text{Fe}]$ and $[\text{Zr}/\text{Fe}]$ ratios are about a factor of 3 to 4 lower than similar metallicity galactic halo stars.

The $[\text{Sr}/\text{Fe}]$ and $[\text{Ba}/\text{Fe}]$ ratios of D119, HD 88609, other Milky Way stars, and the entire sample of literature satellite dwarf galaxy stars published to date are plotted in Figure 2 as a function of $[\text{Fe}/\text{H}]$. The upper limits of these ratios for D119 lie far outside the range of values for Milky Way stars.

D119 is not the only Draco member with a low $[\text{Ba}/\text{Fe}]$ ratio. The star Draco 24, at $[\text{Fe}/\text{H}] = -2.36$, lies about 0.5 dex below Milky Way stars of similar metallicity. More metal-rich stars in Draco show “normal” $[\text{Ba}/\text{Fe}]$ ratios. While it is difficult to prove a trend with just two stars, this may indicate that the whole of Draco started off as Ba-deficient, and

whatever increased the metallicity of the star-forming gas was accompanied by events that created Ba. At high enough metallicity, the initial deficiency became negligible. It would be of interest to determine whether Draco 24 also has a low $[\text{Sr}/\text{Fe}]$ ratio.

The Sextans star, S49 is of similar metallicity to D119 ($[\text{Fe}/\text{H}] = -2.85$), but has a $[\text{Ba}/\text{Fe}]$ value similar to that of Milky Way stars. Whatever caused the lack of Ba (and by extension, Sr) within Draco did not affect a similar star in the Sextans dSph galaxy.

3.1. Comparison to Type II Supernova Models

The production of light elements in Type II supernovae is believed to be a strong function of the progenitor mass. Heavier progenitors should produce more oxygen and magnesium with respect to calcium and titanium. For example, the Type II models of Nomoto et al. (1997), assuming all the ejecta is well-mixed before forming new stars, predict a $[\text{Mg}/\text{Ca}]$ ratio of $+0.47$ for a $70 M_{\odot}$ progenitor and a ratio of -0.42 for a $13 M_{\odot}$ progenitor.

The $[\text{Mg}/\text{Ca}]$ ratio of D119 is $+0.61$, while it is $+0.26$ in HD 88609. The $[\text{Mg}/\text{Fe}]$, $[\text{Ca}/\text{Fe}]$, and $[\text{Mg}/\text{Ca}]$ ratios of these two stars, Milky Way stars, and other dSph stars are shown as a function of $[\text{Fe}/\text{H}]$ in Figure 3. The most metal-poor ($[\text{Fe}/\text{H}] < -2$) dSph stars, including D119, show $[\text{Mg}/\text{Fe}]$ ratios like Milky Way stars, but nearly all the dSph stars show lower $[\text{Ca}/\text{Fe}]$ ratios. However, nearly all the dSph stars show “normal” $[\text{Mg}/\text{Fe}]$ ratios, with D119 being noticeably high. It should be noted that all of the dSph stars, including D119, were analyzed using a nearly identical line list and have similar stellar parameters. This reduces the effects of systematic errors in the relative abundances. The largest uncertainties in the relative abundances come from the line measurement uncertainty in the lower S/N data.

However, within the uncertainties of the measurement, the $[\text{Mg}/\text{Ca}]$ ratio of D119 is consistent with enrichment by only Type II supernovae with progenitor masses of greater than $20 - 25 M_{\odot}$. This would predict $[\text{O}/\text{Fe}]$ and $[\text{Si}/\text{Fe}]$ values for D119 of ≈ 0.5 . Assuming a simple, well-mixed system, the $[\text{Mg}/\text{Ca}]$ ratio in HD 88609 and of all of the other stars in Figure 3 require the addition of ejecta from progenitors of initial lower mass.

In other words, all the stars have similar high-mass contributions as evidenced by their high $[\text{Mg}/\text{Fe}]$ ratios. The difference in $[\text{Mg}/\text{Ca}]$ ratios between the stars therefore depends on the prediction that lower-mass supernovae are significant contributors of Ca. Thus one would predict that D119 would show lower abundances in other elements that are made in significant amounts by low-mass progenitors.

From the Nomoto et al. (1997) models, these elements include Ti, and most of the Fe-group elements (namely, Cr, Mn, Co, Ni, Cu, and Zn). However, interpreting these results is difficult because any combination of the Nomoto et al. (1997) models cannot reproduce the observed abundances of metal-poor stars for several of these elements. For example, all progenitor masses produce sub-solar $[\text{Ti}/\text{Fe}]$ ratios although the vast majority of metal-poor stars show super-solar $[\text{Ti}/\text{Fe}]$ ratios. This problem is a reminder of the speculative nature of using nucleosynthesis models to interpret abundance observations.

Umeda & Nomoto (2002, 2003) explore how progenitor mass, explosion energy and mixing/fallback affect the $[(\text{Zn}, \text{Co}, \text{Mn}, \text{Cr})/\text{Fe}]$ ratios in ejecta from zero-metallicity supernovae. One result is that higher mass progenitors should have lower $[\text{Zn}/\text{Fe}]$ ratios, but mixing/fallback and high explosion energy enhance the $[\text{Zn}/\text{Fe}]$ ratio. While these ratios in D119 may be systematically lower than those seen in HD 88609, the uncertainties are large. An improved analyses with higher S/N spectra could utilize this diagnostic.

Alternatively, the pattern of high Na, Mg, and Al, but low Ca and Ti in D119 is similar to what Chieffi & Limongi (2002) and Imbriani et al. (2001) found for supernova models with high C abundances left by central He burning. However, it is difficult to understand how environment (e.g., in Draco vs. the Milky Way) would affect the central burning properties of massive stars, so this alone could not explain the abundance differences.

A final difficulty in interpreting the nucleosynthesis models is the fate of the massive stars. Heger et al. (2003) find that for the most metal-poor supernovae, those with masses of about 25 to 40 $M_{\odot}$ will leave a black hole formed from material falling back after the explosion. More massive stars (up to the mass range of hypernovae) do not explode at all and collapse directly into black holes. If significant amounts of material are falling back onto the remnant for stars of even 25 $M_{\odot}$, then the interpretation of the results becomes more dependent on the explosion model.

If it can be confirmed that D119 was solely enriched by the ejecta from high-mass supernovae ($M > 20$ to 25 $M_{\odot}$), then this would help eliminate these stars as a site of the r-process. Mathews et al. (1992), Travaglio et al. (1999) and Wanajo et al. (2003) suggest that the r-process primarily takes place in lower mass (8–10 $M_{\odot}$) supernovae. If these theories are correct, D119 may have formed out of the ejecta of higher-mass supernovae before the slower-evolving lower-mass progenitors could evolve.

These results imply the material in D119 came from stars at least as massive as those which Tsujimoto et al. (2000) believe are the primary sites of the r-process. The $[\text{Mg}/\text{H}]$ ratio of D119 is -2.41 , which is right at the point where significant r-process contributions take place in their model. Tsujimoto & Shigeyama (2002) believe that the amount of hydrogen

swept up by supernovae in dSph galaxies should be smaller, so a given supernova shell would have higher metal-to-hydrogen ratios than in the Galactic halo. This would make D119 the product of lower-mass supernova. However, metal-to-metal ratios, such as $[\text{Mg}/\text{Ca}]$, should not be affected by the amount of hydrogen swept up.

3.2. Comparison to Prompt Enrichment Models

The abundance pattern of metal-poor halo stars like HD 88609 are well-fit by the Qian & Wasserburg models. The contribution is dominated by the prompt and low-frequency events, with some high-frequency contributions to produce the Eu seen in these stars. These components cannot reproduce the abundance pattern of D119. In Figure 4(a) we plot the difference in $\log \epsilon$ between the Qian & Wasserburg models.

All of the models overproduce Si, Ca, and Ti. The low-frequency model is included because it is predicted to produce no heavy neutron-capture elements, but the L-model produces no C and too much Cr and Mn. The P-model also produces too much Sr and possibly too much C, O, Y, Zr, and Zn. There is no combination of the two models that can be made to create a satisfactory match to the abundance pattern of D119, mainly due to the overproduction of Sr by the P-model. Only the H-model produces any Ba, so the non-detection of that element eliminates any sizable H-model contribution to D119.

Some of the differences seen in Figure 4(a) may be due to systematic differences (potentially on the order a few tenths of a dex) between the abundance analysis applied here and the previous analyses used to calculate the Qian & Wasserburg models. A systematic analysis might show the abundance ratios of D119 are roughly consistent with the L-model predictions.

If some dSph galaxies did not undergo a prompt enrichment episode, then other very metal-poor stars in these galaxies should show abundance patterns similar to that of D119. If these small galaxies are the CDM building blocks of larger galaxies, then these r-process poor stars should exist around the Milky Way. Therefore, it is of great interest to determine if other dSph galaxies have r-process poor stars and whether there are accreted versions of D119 within the Galaxy.

Zero-metallicity pair-instability hypernovae and normal supernovae have been suggested as the source of any prompt enrichment. Figure 4(b) is a comparison between the Heger & Woosley (2002) hypernovae models and D119, plotting the scaled models for He core masses of 75, 100, and 125 $M_{\odot}$. One characteristic of the nucleosynthesis of pair production supernovae is the pronounced odd-even effect. This pattern would make it difficult for any

combination of models to reproduce the abundances of D119. For example, the models all overproduce Cr and underproduce Sc. The zero-metallicity models of both Umeda & Nomoto (2002) for both supernovae (13 to 30 $M_{\odot}$) and hypernovae (150 to 270 $M_{\odot}$) also show this same strong odd-even effect.

4. Another Low [Ba/Fe] Star: BD +80 245

The unusual α -poor nature of the metal-poor halo star BD +80 245 was discovered by Carney et al. (1997). Unlike most metal-poor stars at $[\text{Fe}/\text{H}] \approx -2$, which have super-solar $[\alpha/\text{Fe}]$ ratios (see Figure 3), BD +80 245 shows sub-solar ratios. Further, Fulbright (2000) found that $[\text{Ba}/\text{Fe}]$ and $[\text{Eu}/\text{Fe}]$ are ~ 1 and ~ 2 dex below the ratios seen in other halo stars of similar metallicity.

There are two possible explanations for the abundance pattern in BD +80 245. Since low-mass Type II supernovae produce lower α/Fe ratios, the star could be the product of a system only contaminated by low-mass events. However, the evidence provided here by D119 points to low-mass events being producers of r-process elements, so BD +80 245 should show high $[\text{Ba}/\text{Fe}]$ and $[\text{Eu}/\text{Fe}]$ ratios, in contrast to observations. The alternative explanation, that BD +80 245 is formed out of some Type II ejecta mixed with Type Ia ejecta, is a more likely explanation for the abundance pattern found in this star. Ivans et al. (2003) further explores the possible enrichment history of this class of metal-poor, α -poor stars.

5. Summary

We have analyzed new high S/N spectra of the metal-poor Draco dSph star D119, confirming ($[\text{Fe}/\text{H}] = -2.96$) that this star remains the most metal poor star presently known in any dwarf galaxy. Our spectra of D119 show *no measurable lines for any neutron capture elements*, a phenomenon not previously noted in any star. One other star nearly as metal poor in the Sextans dwarf spheroidal does not have the peculiar composition of D119. While binary mass transfer processes can account for excesses of neutron capture elements, we can imagine no astrophysical process which might deplete so completely the heavy elements. The unique composition of D119 must be attributed to the chemical enrichment history of the gas from which it was formed.

The abundance ratios of Draco 119 are consistent with that expected for material formed from the ejecta of massive Type II supernovae, a hypothesis that would also be consistent with the great age and low metallicity of the Draco dwarf spheroidal galaxy Grillmair et

al. (1998). This is consistent with the predictions of Mathews et al. (1992), Travaglio et al. (1999) and Wanajo et al. (2003), which places the r-process in low mass Type II supernovae, and may conflict with the prediction of Tsujimoto et al. (2000) who propose intermediate to massive Type II SNe as the site of the the r-process. The heavy element abundance distribution disagrees with the prompt enrichment predictions of Qian & Wasserburg and the pair-production supernovae models of Heger & Woosley (2002) and Umeda & Nomoto (2002). At present we find no supernova model with predicted yields that can fit our observations.

The authors would like to thank J. Johnson and M. Shetrone for granting us access to their spectra. We would also like to thank the referees, S. Ryan and M. Shetrone, plus G. Wasserburg, K. Nomoto, C. Sneden, Wal Sargent and Brad Hansen for their many valuable suggestions. We are especially grateful to the staff of Keck Observatory for their assistance, and S. Vogt and his team for the creation of HIRES. JPF would like to thank Andy McWilliam for many useful conversations. RMR acknowledges support from grant AST-0098739 from the National Science Foundation. The authors acknowledge the cultural role that the summit of Mauna Kea has had within the indigenous Hawaiian community. We are fortunate to have the opportunity to conduct observations from this mountain.

REFERENCES

- Alonso, A., Arribas, S. & Martinez-Roger, C. 1999a, A&AS 140, 261
- Alonso, A., Arribas, S. & Martinez-Roger, C. 1999b, A&AS 139, 335
- Bonifacio, P., Hill, V., Molaro, P., Pasquini, L., Di Marcantonio, P. & Santin, P. 2000, A&A, 359, 663
- Brown, J. 1987, ApJ, 317, 701
- Burbidge, E. M., Burbidge, G. R., Fowler, W. A. & Hoyle, F. 1957, RvMP, 29, 547
- Carney, B. W., Wright, J. S., Sneden, C., Laird, J. B., Aguilar, L. A., & Latham, D. W. 1997, AJ, 114, 363
- Cayrel, R. 1988, in IAU Symp. 132, The Impact of Very High S/N Spectroscopy on Stellar Physics, ed. G. Cayrel de Strobel & Monique Spite (Kluwer: Dordrecht), 345
- Chieffi, A. & Limongi, M. 2002, ApJ, 577, 281
- Fulbright, J. P. 2000, AJ, 120, 1841

- Grillmair, C.J., Mould, J.R., Holtzman, J.A., Worthey, G., Ballester, G.E., Clarke, J.T., Crisp, D., Evans, R.W., Gallagher, J.S., III, & Burrows, C.J. 1998, *AJ*, 115, 144
- Hanson, R. B., Sneden, C., Kraft, R. P., Fulbright, J. 1998, *AJ*, 116, 128
- Heger, A & Woosley, S. E. 2002, *ApJ*, 567, 532
- Heger, A., Fryer, C. L., Woosley, S. E., Langer, N. & Hartmann, D. H. 2003, *ApJ*, 591, 288
- Ivans, I. I., Sneden, C., Renee James, C., Preston, G., Fulbright, J. P., Hoflick, P. A., Carney, B. & Wheeler, J. C. 2003, accepted
- Johnson, J. A. 2002, *ApJS*, 139, 219
- Johnson, J. A. & Bolte, M. 2002, *ApJ*, 579, 616
- Kraft, R. P. & Ivans, I. I. 2003, in publication
- Imbriani, G., Limongi, M., Gialanella, L., Terrasi, F., Straniero, O., & Chieffi, A. 2001, *ApJ*, 558, 903
- Mateo, M. 1998, *ARAA*, 36, 435
- Mathews, G. J., Bazan, G. & Cowan, J. J. 1992, *ApJ*, 391, 719
- McWilliam, A., Preston, G., Sneden, C. & Searle, L. 1995, *AJ*, 109, 2757
- McWilliam, A. 1998, *AJ*, 115, 1640
- McWilliam, A., Rich, R. M. & Smecker-Hane, T. A. 2003, *ApJ*, 593, L145
- Moore, C. 1966, *The Solar Spectrum 2935 Å to 8770 Å*, NBSM Monograph 61
- Nakamura, T., Umeda, H., Nomoto, K., Thieleman, F.-K. & Burrows, A. 1999, *ApJ*, 517, 193
- Nomoto, K., Hashimoto, M., Tsujimoto, T., Thielemann, F.-K. & Kubo, Y. 1997, *Nuc Phys A*, 616, 79
- Qian, Y.-Z. 2003, astro-ph/0301422
- Qian, Y.-Z. & Wasserburg, G. J. 2001, *ApJ*, 559, 925
- Qian, Y.-Z. & Wasserburg, G. J. 2002, *ApJ*, 567, 515
- Ryan, S. G., Norris, J. E. & Beers, T. C. 1996, *ApJ*, 471, 254

- Shetrone, M. D., Bolte, M. & Stetson, P. B. 1998, *AJ*, 115, 1888
- Shetrone, M. D., Côte, P. & Sargent, W. L. W., 2001, *ApJ*, 548, 592
- Shetrone, M. D., Venn, K. A., Tolstoy, E. Primas, F., Hill, V. & Kaufer, A. 2003, *A&A*, 125, 707
- Snedden, C. 1973, PhD Dissertation, Univ. of Texas at Austin
- Snedden, C. & Cowan, J. J. 2003, *Science*, 299, 70
- Stetson, P. B. 1979, *AJ*, 84, 1149
- Travaglio, C., Galli, D., Gallino, R., Busso, M., Ferrini, F. & Straniero, O. 1999, *ApJ*, 521, 691
- Truran, J. W., Cowan, J. J., Pilachowski, C. A. & Sneden, C. 2002, *PASP*, 114, 1293
- Tsujimoto, T., Shigeyama, T. & Yoshii, Y. 2000, *ApJL*, 531, L33
- Tsujimoto, T. & Shigeyama, T. 2002, *ApJL*, 571, L93
- Umeda, H. & Nomoto, K. 2002, *ApJ*, 565, 385
- Umeda, H. & Nomoto, K. 2003, *Nature*, 422, 871
- Vogt, S. S. et al. 1994, *SPIE*, 2198, 362
- Wanajo, S., Tamamura, M., Itoh, N., Nomoto, K., Ishimaru, Y., Beers, T. C., Satoshi, N. 2003, *ApJ*, 593, 968

Table 1. Line List

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
4288.74	CH ^a	0.64	−1.138	17	19
4288.74	CH	0.64	−1.115	bl	bl
4307.31	CH	0.16	−1.475	29	22
4310.09	CH	0.10	−1.534	28	40
4310.11	CH	0.10	−1.595	bl	bl
4313.59	CH	0.02	−1.923	30	21
4313.65	CH	0.02	−1.923	bl	bl
6300.31	[O I]	0.00	−9.750	< 8	...
5889.97	Na I	0.00	hfs ^b	190	180
4703.00	Mg I	4.34	−0.520	56	61
5172.70	Mg I	2.71	−0.381	192	202
5183.62	Mg I	2.72	−0.170	219	214
5528.42	Mg I	4.34	−0.500	68	64
3961.54	Al I	0.00	hfs	140	134
4102.94	Si I	1.91	−3.140	< 51	75
5684.52	Si I	4.93	−1.650	< 10	7
5708.41	Si I	4.95	−1.470	< 10	6
4318.65	Ca I	1.90	−0.210	21	44
6122.23	Ca I	1.89	−0.320	45	59
6439.08	Ca I	2.52	0.390	31	52
4320.75	Sc II	0.61	−0.250	97	82
4354.61	Sc II	0.61	−1.580	25	22
4670.41	Sc II	1.36	−0.580	25	24
5239.82	Sc II	1.45	−0.760	22	13
5526.82	Sc II	1.77	+0.020	46	30
4533.24	Ti I	0.85	+0.540	27	43
4534.78	Ti I	0.84	+0.340	34	34
4555.49	Ti I	0.85	−0.430	15	9
4981.73	Ti I	0.85	+0.560	49	22

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
4999.50	Ti I	0.83	+0.310	29	39
5014.24	Ti I	0.81	+0.110	48	...
5020.03	Ti I	0.84	−0.350	10	12
5022.87	Ti I	0.83	−0.370	19	...
5039.96	Ti I	0.21	−1.070	29	24
5064.65	Ti I	0.05	−0.930	41	30
5173.74	Ti I	0.00	−1.060	22	26
5192.97	Ti I	0.02	−0.950	34	32
5210.39	Ti I	0.05	−0.820	24	34
4394.06	Ti II	1.22	−1.770	52	54
4395.04	Ti II	1.08	−0.510	121	137
4395.84	Ti II	1.24	−1.970	39	41
4398.29	Ti II	1.22	−2.780	13	11
4443.80	Ti II	1.08	−0.700	110	116
4444.56	Ti II	1.12	−2.210	32	38
4450.48	Ti II	1.08	−1.510	68	81
4493.52	Ti II	1.08	−2.830	11	14
4501.27	Ti II	1.12	−0.760	117	116
4708.66	Ti II	1.24	−2.370	14	26
4798.53	Ti II	1.08	−2.670	24	26
5336.79	Ti II	1.58	−1.630	42	43
5129.16	Ti II	1.89	−1.390	27	...
5154.07	Ti II	1.57	−1.920	16	...
5226.55	Ti II	1.57	−1.300	70	...
5381.01	Ti II	1.57	−2.080	23	...
4379.23	V I	0.30	+0.550	44	20
4389.98	V I	0.28	+0.270	18	11
4254.35	Cr I	0.00	−0.110	107	109
4616.13	Cr I	0.98	−1.200	12	13

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
4646.17	Cr I	1.03	−0.720	20	26
5296.70	Cr I	0.98	−1.400	15	12
5345.81	Cr I	1.00	−0.980	19	21
5409.80	Cr I	1.03	−0.720	31	33
4652.17	Cr I	1.00	−1.030	23	17
5206.04	Cr I	0.94	+0.019	68	...
5409.80	Cr I	1.03	−0.720	36	33
4030.76	Mn I	0.00	hfs	96	117
4033.07	Mn I	0.00	hfs	89	109
4034.49	Mn I	0.00	hfs	106	97
4250.13	Fe I	2.47	−0.370	68	84
4260.47	Fe I	2.40	+0.140	92	108
4282.41	Fe I	2.17	−0.780	83	82
4337.05	Fe I	1.56	−1.660	84	89
4375.93	Fe I	0.00	−2.990	126	126
4404.75	Fe I	1.56	−0.100	150	149
4415.13	Fe I	1.61	−0.620	128	128
4427.31	Fe I	0.05	−3.000	140	...
4430.62	Fe I	2.22	−1.620	48	...
4443.20	Fe I	2.86	−1.040	33	27
4445.48	Fe I	0.09	−5.400	12	10
4447.73	Fe I	2.22	−1.340	66	58
4461.65	Fe I	0.09	−3.170	127	122
4489.74	Fe I	0.12	−3.930	86	77
4494.57	Fe I	2.20	−1.100	53	76
4531.15	Fe I	1.49	−2.110	80	75
4602.94	Fe I	1.49	−2.180	83	14
4630.13	Fe I	2.28	−2.590	19	9
4632.92	Fe I	1.61	−2.900	41	24

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
4647.43	Fe I	2.94	−1.350	31	16
4733.60	Fe I	1.49	−2.950	37	31
4736.77	Fe I	3.20	−0.750	37	30
4859.74	Fe I	2.86	−0.760	59	...
4871.32	Fe I	2.85	−0.360	79	64
4872.14	Fe I	2.87	−0.570	72	52
4890.75	Fe I	2.86	−0.390	74	62
4891.49	Fe I	2.84	−0.110	76	76
4903.31	Fe I	2.87	−0.930	39	38
4918.99	Fe I	2.85	−0.340	84	65
4920.50	Fe I	2.82	+0.070	83	89
4924.77	Fe I	2.28	−2.250	24	19
4938.81	Fe I	2.86	−1.080	32	28
4939.69	Fe I	0.86	−3.300	72	58
4994.13	Fe I	0.92	−3.040	76	70
5006.12	Fe I	2.83	−0.660	70	59
5041.07	Fe I	0.95	−3.090	78	71
5049.82	Fe I	2.28	−1.340	68	57
5051.64	Fe I	0.92	−2.760	92	90
5068.77	Fe I	2.93	−1.040	29	26
5079.23	Fe I	2.20	−2.030	31	34
5079.74	Fe I	0.99	−3.180	71	58
5083.34	Fe I	0.96	−2.920	86	77
5123.72	Fe I	1.01	−3.030	77	65
5127.36	Fe I	0.92	−3.270	69	61
5141.73	Fe I	2.42	−1.960	14	16
5142.93	Fe I	0.96	−3.080	65	69
5151.92	Fe I	1.01	−3.280	63	47
5166.29	Fe I	0.00	−4.160	100	84

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
5171.60	Fe I	1.49	−1.750	109	97
5191.45	Fe I	3.03	−0.550	52	44
5192.34	Fe I	2.99	−0.420	54	54
5194.94	Fe I	1.56	−2.050	87	79
5198.71	Fe I	2.22	−2.090	24	25
5216.28	Fe I	1.61	−2.110	78	71
5217.39	Fe I	3.21	−1.070	25	14
5225.53	Fe I	0.11	−4.750	55	35
5232.94	Fe I	2.94	−0.100	83	77
5247.05	Fe I	0.09	−4.910	38	26
5250.21	Fe I	0.12	−4.900	40	28
5269.54	Fe I	0.86	−1.330	173	161
5281.79	Fe I	3.03	−0.830	34	30
5283.62	Fe I	3.23	−0.520	47	39
5302.30	Fe I	3.28	−0.720	28	26
5307.37	Fe I	1.61	−2.950	34	24
5324.19	Fe I	3.21	−0.100	55	...
5332.90	Fe I	1.55	−2.780	50	34
5339.93	Fe I	3.27	−0.650	27	27
5341.02	Fe I	1.60	−1.950	98	83
5371.50	Fe I	0.96	−1.644	157	141
5383.37	Fe I	4.31	+0.640	33	24
5393.17	Fe I	3.24	−0.710	28	25
5397.13	Fe I	0.92	−1.950	158	135
5404.15	Fe I	4.42	+0.520	20	27
5405.78	Fe I	0.99	−1.800	144	132
5415.20	Fe I	4.37	+0.640	26	20
5429.70	Fe I	0.95	−1.880	150	134
5434.53	Fe I	1.01	−2.080	126	...

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
5501.46	Fe I	0.95	−3.050	85	77
5506.78	Fe I	0.99	−2.700	99	89
5615.66	Fe I	3.33	+0.050	71	...
6136.62	Fe I	2.45	−1.400	56	52
6137.70	Fe I	2.59	−1.366	59	43
6191.57	Fe I	2.43	−1.416	70	...
6219.29	Fe I	2.20	−2.438	37	19
6230.74	Fe I	2.56	−1.276	73	...
6252.57	Fe I	2.40	−1.757	57	42
6421.36	Fe I	2.28	−2.014	30	35
6430.86	Fe I	2.18	−1.946	45	45
4508.29	Fe II	2.86	−2.330	40	42
4515.34	Fe II	2.84	−2.480	24	31
4555.89	Fe II	2.83	−2.390	40	39
5018.45	Fe II	2.89	−1.220	101	106
5197.56	Fe II	3.23	−2.100	35	27
5234.62	Fe II	3.22	−2.230	27	36
5276.00	Fe II	3.20	−1.940	41	38
4121.32	Co I	0.92	hfs	82	77
5035.36	Ni I	3.63	+0.290	14	12
5137.08	Ni I	1.68	−1.990	13	20
5476.91	Ni I	1.83	−0.890	65	64
5105.54	Cu I	1.39	−1.720	< 9	...
4810.55	Zn I	4.08	−0.170	< 10	15
4077.71	Sr II	0.00	hfs	< 53	169
4215.52	Sr II	0.00	hfs	< 25	160
4883.69	Y II	1.08	+0.070	< 10	20
4900.11	Y II	1.03	−0.090	< 10	21
5087.42	Y II	1.08	−0.170	< 9	12

Table 1—Continued

Wavelength Å	Ion	E.P. eV	log gf	D119 EW mÅ	HD 88609 EW mÅ
4208.98	Zr II	0.71	−0.460	< 25	30
4554.03	Ba II	0.00	hfs	< 14	91
4934.08	Ba II	0.00	hfs	< 10	...
4061.09	Nd II	0.47	+0.300	< 35	5
4109.46	Nd II	0.32	+0.180	< 35	5
4129.72	Eu II	0.00	hfs	< 41	9

^aCH line analysis assumed a disassociation energy of 3.47 eV. Lines with “bl” in the EW columns were lines analyzed as blends with the line preceeding it in the list, with the given EW assumed to cover both lines.

^bines with log gf values marked as “hfs” were treated as blend of many hyperfine components. Hyperfine data taken from Johnson (2002), McWilliam (1998), and McWilliam et al. (1995).

Table 2. Observed Abundances

Species	Draco 119			HD 88609		
	$\log \epsilon(X)$	$[X/Fe]^a$	σ	$\log \epsilon(X)$	$[X/Fe]^a$	σ
C ^b	5.18	−0.48	0.26	5.19	−0.43	0.14
[O I]	< 6.6	< 0.8	...	...	...	...
Na I	3.74	+0.36	0.12	3.53	+0.14	0.07
Mg I	5.17	+0.54	0.22	5.23	+0.59	0.07
Al I	3.28	−0.24	0.49	3.17	−0.36	0.10
Si I	< 4.8	< +0.2	...	5.18	+0.57	0.11
Ca I	3.34	−0.07	0.17	3.75	+0.33	0.08
Sc II	0.43	+0.30	0.18	0.40	+0.13	0.07
Ti I	2.11	+0.07	0.13	2.22	+0.17	0.05
Ti II	2.23	+0.21	0.15	2.48	+0.32	0.06
V I	1.19	+0.14	0.29	0.94	−0.12	0.05
Cr I	2.30	−0.42	0.14	2.45	−0.28	0.05
Mn I	1.75	−0.69	0.48	2.15	−0.30	0.11
Fe I	4.57	−2.95	0.20	4.58	−2.94	0.14
Fe II	4.55	−2.97	0.12	4.69	−2.83	0.05
Co I	< 2.1	< +0.1	...	2.10	+0.12	0.07
Ni I	3.15	−0.15	0.17	3.31	+0.00	0.06
Cu I	< 1.17	< −0.1	...	...	...	...
Zn I	< 1.7	< +0.1	...	2.01	+0.35	0.13
Sr II	< −2.6	< −2.5	...	0.00	−0.07	0.12
Y II	< −1.3	< −0.5	...	−0.80	−0.21	0.07
Zr II	< −0.2	< +0.1	...	0.04	+0.27	0.08
Ba II	< −3.3	< −2.6	...	−1.61	−0.91	0.10
Nd II	< −0.7	< +0.8	...	−1.62	−0.27	0.14
Eu II	< −2.0	< +0.4	...	−2.59	−0.50	0.08

^a[Fe/H] ratios given for Fe I and Fe II.

^bBased on G-band CH lines.

Fig. 1.— Two sample spectral regions showing the difference in neutron-capture line strengths between D119 (solid) and the halo star HD 88609 (dotted), which has the same metallicity as D119. The left panel shows the Sr II 4215 Å line for both stars. Note the near identical strength of the Fe I line. The right panel shows the Ba II 4554 Å line for both stars.

Fig. 2.— The [Sr/Fe] and [Ba/Fe] ratios of D119, HD 88609, field halo stars, and dSph giants plotted against [Fe/H]. The field star data are from McWilliam (1995), Ryan et al. (1996), Burris et al. (2000), Fulbright (2000), Johnson (2002) and Ivans et al. (2003). All of the field stars (with the exception of BD +80+245) are plotted using the same symbol to prevent confusion. Except for D119, the dSph sample comes from Shetrone et al. (2001) and Shetrone et al. (2003). In both plots, the upper limits for D119 clearly lie below other stars of similar metallicity.

Fig. 3.— The [Mg/Fe], [Ca/Fe] and [Mg/Ca] ratios of D119 and the other stars plotted in Figure 2 (same symbols used). In the top panel, D119 and the other very metal-poor dSph stars show [Mg/Fe] ratios similar to that seen in the Milky Way population. The dSph stars, however, show lower [Ca/Fe] ratios in the middle panel. In the bottom panel, D119 show a very high [Mg/Ca] ratio, which is the result of having a slightly higher [Mg/Fe] ratio and slightly lower [Ca/Fe] ratio than similar stars (such as Sextans S49). Overall, the dSph stars show [Mg/Ca] ratios not too different than the Milky Way sample, suggesting that with the exception of D119, the mass function of Type II supernovae progenitors that polluted the two groups was similar.

Fig. 4.— Comparison of theoretical models to the observed abundances of D119. In both panels, the quantity $\Delta \log \epsilon$ is defined as $\log \epsilon(\text{Model}) - \log \epsilon(\text{D119})$, and the models are scaled to the Fe abundance of D119. Positive values indicate overproduction by the models, while the zero-line indicates a perfect match to D119. The upward-pointing arrows denote elements for which the marked points are lower limits based on the adopted upper limit of the abundance measured in D119. (a) Comparison of the P- and L-model predictions of Qian & Wasserburg to the observed abundances of D119. The prediction of the carbon abundance from the P-model is an upper limit. (b) Comparison of the Heger & Woosley (2002) predictions of the nucleosynthesis of pair instability supernovae to the observed abundances of D119. The three models are based on He-core masses of 75, 100 and 125 $M_{\odot}$.







